

8 Matrix Multiplication

1. The three-post snap group can be represented by a set of graphs, each with three towns. The posts are the towns and the elastic bands are the roads. For example,

$$A = \begin{matrix} & \text{to} \\ & 1 & 2 & 3 \\ \text{from} & \begin{bmatrix} 1 & 0 & 0 \\ 2 & 0 & 1 \\ 3 & 0 & 1 \end{bmatrix} \end{matrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} & \textcircled{3} \\ \hline & \textcircled{2} \\ \hline \end{array}$$

- (a) Draw the graphs and transportation matrices for this group.

Here they are!

$$\begin{array}{l} I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} & \textcircled{3} \\ \hline & \textcircled{2} \\ \hline \end{array} \\ B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} \leftrightarrow \textcircled{3} \\ \hline & \textcircled{2} \\ \hline \end{array} \\ D = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} \rightarrow \textcircled{3} \\ \hline \textcircled{1} \rightarrow \textcircled{2} \\ \hline \textcircled{2} \rightarrow \textcircled{3} \\ \hline \end{array} \\ A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} & \textcircled{3} \\ \hline & \textcircled{2} \\ \hline \end{array} \\ C = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} \rightarrow \textcircled{2} \\ \hline & \textcircled{3} \\ \hline \end{array} \\ E = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \leftrightarrow \begin{array}{|c|} \hline \textcircled{1} \leftarrow \textcircled{3} \\ \hline \textcircled{1} \rightarrow \textcircled{2} \\ \hline \textcircled{2} \rightarrow \textcircled{3} \\ \hline \end{array} \end{array}$$

- (b) Try a few multiplications and notice the isomorphism to the snap group.

Before, we found that $A \cdot B = E$. But does this work with the matrices? We have

$$\begin{aligned} AB &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} \langle 1, 0, 0 \rangle \cdot \langle 0, 0, 1 \rangle & \langle 1, 0, 0 \rangle \cdot \langle 0, 1, 0 \rangle & \langle 1, 0, 0 \rangle \cdot \langle 1, 0, 0 \rangle \\ \langle 0, 0, 1 \rangle \cdot \langle 0, 0, 1 \rangle & \langle 0, 0, 1 \rangle \cdot \langle 0, 1, 0 \rangle & \langle 0, 0, 1 \rangle \cdot \langle 1, 0, 0 \rangle \\ \langle 0, 1, 0 \rangle \cdot \langle 0, 0, 1 \rangle & \langle 0, 1, 0 \rangle \cdot \langle 0, 1, 0 \rangle & \langle 0, 1, 0 \rangle \cdot \langle 1, 0, 0 \rangle \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} = D. \end{aligned}$$

Huh?

The issue is simple. Matrix multiplication, just like the snap operation, is not commutative, and we need to flip the order of the matrices so it represents taking B first, then A . After all, that's what we defined $A \cdot B$ to be.

$$\begin{aligned}
BA &= \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \\
&= \begin{bmatrix} \langle 0,0,1 \rangle \cdot \langle 1,0,0 \rangle & \langle 0,0,1 \rangle \cdot \langle 0,0,1 \rangle & \langle 0,0,1 \rangle \cdot \langle 0,1,0 \rangle \\ \langle 0,1,0 \rangle \cdot \langle 1,0,0 \rangle & \langle 0,1,0 \rangle \cdot \langle 0,0,1 \rangle & \langle 0,1,0 \rangle \cdot \langle 0,1,0 \rangle \\ \langle 1,0,0 \rangle \cdot \langle 1,0,0 \rangle & \langle 1,0,0 \rangle \cdot \langle 0,0,1 \rangle & \langle 1,0,0 \rangle \cdot \langle 0,1,0 \rangle \end{bmatrix} \\
&= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} = E.
\end{aligned}$$

This works for any of the matrices.

2. Using 3×3 matrices A and B from this section, compute

For reference, the matrices are

$$A = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 1 \\ 2 & 0 & 1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}.$$

(a) $AA = A^2$

We use the column vector/row vector approach.

$$\begin{aligned}
AA &= \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 1 \\ 2 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 1 \\ 2 & 0 & 1 & 1 \end{bmatrix} \\
&= \begin{bmatrix} \langle 1,1,2,2 \rangle \cdot \langle 1,1,2,2 \rangle & \langle 1,1,2,2 \rangle \cdot \langle 1,1,1,0 \rangle & \langle 1,1,2,2 \rangle \cdot \langle 2,1,1,1 \rangle & \langle 1,1,2,2 \rangle \cdot \langle 2,0,1,1 \rangle \\ \langle 1,1,1,0 \rangle \cdot \langle 1,1,2,2 \rangle & \langle 1,1,1,0 \rangle \cdot \langle 1,1,1,0 \rangle & \langle 1,1,1,0 \rangle \cdot \langle 2,1,1,1 \rangle & \langle 1,1,1,0 \rangle \cdot \langle 2,0,1,1 \rangle \\ \langle 2,1,1,1 \rangle \cdot \langle 1,1,2,2 \rangle & \langle 2,1,1,1 \rangle \cdot \langle 1,1,1,0 \rangle & \langle 2,1,1,1 \rangle \cdot \langle 2,1,1,1 \rangle & \langle 2,1,1,1 \rangle \cdot \langle 2,0,1,1 \rangle \\ \langle 2,0,1,1 \rangle \cdot \langle 1,1,2,2 \rangle & \langle 2,0,1,1 \rangle \cdot \langle 1,1,1,0 \rangle & \langle 2,0,1,1 \rangle \cdot \langle 2,1,1,1 \rangle & \langle 2,0,1,1 \rangle \cdot \langle 2,0,1,1 \rangle \end{bmatrix} \\
&= \begin{bmatrix} 10 & 4 & 7 & 6 \\ 4 & 3 & 4 & 3 \\ 7 & 4 & 7 & 6 \\ 6 & 3 & 6 & 6 \end{bmatrix}.
\end{aligned}$$

(b) AB

$$\begin{aligned}
AB &= \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 1 \\ 2 & 0 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix} \\
&= \begin{bmatrix} \langle 1,1,2,2 \rangle \cdot \langle 1,1,0,0 \rangle & \langle 1,1,2,2 \rangle \cdot \langle 0,1,1,0 \rangle & \langle 1,1,2,2 \rangle \cdot \langle 0,0,1,1 \rangle & \langle 1,1,2,2 \rangle \cdot \langle 1,0,0,1 \rangle \\ \langle 1,1,1,0 \rangle \cdot \langle 1,1,0,0 \rangle & \langle 1,1,1,0 \rangle \cdot \langle 0,1,1,0 \rangle & \langle 1,1,1,0 \rangle \cdot \langle 0,0,1,1 \rangle & \langle 1,1,1,0 \rangle \cdot \langle 1,0,0,1 \rangle \\ \langle 2,1,1,1 \rangle \cdot \langle 1,1,0,0 \rangle & \langle 2,1,1,1 \rangle \cdot \langle 0,1,1,0 \rangle & \langle 2,1,1,1 \rangle \cdot \langle 0,0,1,1 \rangle & \langle 2,1,1,1 \rangle \cdot \langle 1,0,0,1 \rangle \\ \langle 2,0,1,1 \rangle \cdot \langle 1,1,0,0 \rangle & \langle 2,0,1,1 \rangle \cdot \langle 0,1,1,0 \rangle & \langle 2,0,1,1 \rangle \cdot \langle 0,0,1,1 \rangle & \langle 2,0,1,1 \rangle \cdot \langle 1,0,0,1 \rangle \end{bmatrix} \\
&= \begin{bmatrix} 3 & 2 & 3 & 4 \\ 1 & 2 & 2 & 1 \\ 3 & 3 & 2 & 2 \\ 3 & 2 & 1 & 2 \end{bmatrix}.
\end{aligned}$$

(c) BA

$$\begin{aligned}
 BA &= \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 & 2 \\ 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 1 \\ 2 & 0 & 1 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} \langle 1, 1, 0, 0 \rangle \cdot \langle 1, 1, 2, 2 \rangle & \langle 1, 1, 0, 0 \rangle \cdot \langle 1, 1, 1, 0 \rangle & \langle 1, 1, 0, 0 \rangle \cdot \langle 2, 1, 1, 1 \rangle & \langle 1, 1, 0, 0 \rangle \cdot \langle 2, 0, 1, 1 \rangle \\ \langle 0, 1, 1, 0 \rangle \cdot \langle 1, 1, 2, 2 \rangle & \langle 0, 1, 1, 0 \rangle \cdot \langle 1, 1, 1, 0 \rangle & \langle 0, 1, 1, 0 \rangle \cdot \langle 2, 1, 1, 1 \rangle & \langle 0, 1, 1, 0 \rangle \cdot \langle 2, 0, 1, 1 \rangle \\ \langle 0, 0, 1, 1 \rangle \cdot \langle 1, 1, 2, 2 \rangle & \langle 0, 0, 1, 1 \rangle \cdot \langle 1, 1, 1, 0 \rangle & \langle 0, 0, 1, 1 \rangle \cdot \langle 2, 1, 1, 1 \rangle & \langle 0, 0, 1, 1 \rangle \cdot \langle 2, 0, 1, 1 \rangle \\ \langle 1, 0, 0, 1 \rangle \cdot \langle 1, 1, 2, 2 \rangle & \langle 1, 0, 0, 1 \rangle \cdot \langle 1, 1, 1, 0 \rangle & \langle 1, 0, 0, 1 \rangle \cdot \langle 2, 1, 1, 1 \rangle & \langle 1, 0, 0, 1 \rangle \cdot \langle 2, 0, 1, 1 \rangle \end{bmatrix} \\
 &= \begin{bmatrix} 2 & 2 & 3 & 2 \\ 3 & 2 & 2 & 1 \\ 4 & 1 & 2 & 2 \\ 3 & 1 & 3 & 3 \end{bmatrix}.
 \end{aligned}$$


(d) B^2

$$\begin{aligned}
 BA &= \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} \langle 1, 1, 0, 0 \rangle \cdot \langle 1, 0, 0, 1 \rangle & \langle 1, 1, 0, 0 \rangle \cdot \langle 1, 1, 0, 0 \rangle & \langle 1, 1, 0, 0 \rangle \cdot \langle 0, 1, 1, 0 \rangle & \langle 1, 1, 0, 0 \rangle \cdot \langle 0, 0, 1, 1 \rangle \\ \langle 0, 1, 1, 0 \rangle \cdot \langle 1, 0, 0, 1 \rangle & \langle 0, 1, 1, 0 \rangle \cdot \langle 1, 1, 0, 0 \rangle & \langle 0, 1, 1, 0 \rangle \cdot \langle 0, 1, 1, 0 \rangle & \langle 0, 1, 1, 0 \rangle \cdot \langle 0, 0, 1, 1 \rangle \\ \langle 0, 0, 1, 1 \rangle \cdot \langle 1, 0, 0, 1 \rangle & \langle 0, 0, 1, 1 \rangle \cdot \langle 1, 1, 0, 0 \rangle & \langle 0, 0, 1, 1 \rangle \cdot \langle 0, 1, 1, 0 \rangle & \langle 0, 0, 1, 1 \rangle \cdot \langle 0, 0, 1, 1 \rangle \\ \langle 1, 0, 0, 1 \rangle \cdot \langle 1, 0, 0, 1 \rangle & \langle 1, 0, 0, 1 \rangle \cdot \langle 1, 1, 0, 0 \rangle & \langle 1, 0, 0, 1 \rangle \cdot \langle 0, 1, 1, 0 \rangle & \langle 1, 0, 0, 1 \rangle \cdot \langle 0, 0, 1, 1 \rangle \end{bmatrix} \\
 &= \begin{bmatrix} 1 & 2 & 1 & 0 \\ 0 & 1 & 2 & 1 \\ 1 & 0 & 1 & 2 \\ 2 & 1 & 0 & 1 \end{bmatrix}.
 \end{aligned}$$

(e) Which one (AB and BA) represents taking a step by walking, then by bus?

Since A is walking and B is bus, we know that AB is a step by walking, then by bus. Unlike most of the operations we've been doing, it is not right-to-left!

One way to conceptualize this is to draw an arrow from "from" to "to" for each matrix, then join the arrows. For example, for the product AB , it should look like this:



This arrow shows the order in which paths are taken.

(f) Use your calculator to check your computations of A^2 , AB , BA , and B^2 .

Here's some instructions on multiplying matrices on various TI calculators:

TI-83/TI-84: Press "2nd" and " x^{-1} ," or if your calculator has it, the "MATRIX" button, to enter the matrix editing page. Navigate to the "EDIT" menu, then navigate to the desired name for the first matrix. Press enter to select that matrix, then type in the size and values of the matrix. Repeat this for the second matrix. When you're ready to multiply them, press "2nd" and " x^{-1} " again, but stay in the "NAMES" menu. Navigate to the

first matrix to multiply, and press enter. Repeat this for the second matrix. Finally, pressing enter to calculate will give us the result of the multiplication (or an error if the dimensions are incorrect).

TI Nspire: Press the button to the bottom left of " \leftarrow^{del} " that looks like " $\left[\begin{array}{c} \blacksquare \\ \blacksquare \\ \blacksquare \end{array} \right]$ ". Navigate to the button that looks like a blank 3×3 matrix and press enter. Enter in the size of the first matrix, then the values. Repeat this process for the second matrix, then multiply the two matrices by pressing enter.

Assuming I didn't make a large oopsie, those answers are all correct. :P

3. Write a 3×3 matrix T that shows the following scenario: you can go from town B to C , C to D , and D to B by train, in exactly one way each, and not backwards.

$$T = \begin{array}{c} \text{to} \\ B \quad C \quad D \\ \text{from} \\ \begin{array}{c} B \\ C \\ D \end{array} \end{array} \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}.$$

- (a) Why can't you add this matrix to matrices A or B ?

They have different dimensions!

- (b) Rewrite matrix T so that it *can* be meaningfully added to matrices A and B . What did you do to its dimensions?

We need T to be 4×4 , and we want the entries A, B, C, D to line up properly. Thus, we insert 0s in the A column and A row, as shown:

$$T = \begin{array}{c} \text{to} \\ A \quad B \quad C \quad D \\ \text{from} \\ \begin{array}{c} A \\ B \\ C \\ D \end{array} \end{array} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \end{bmatrix}.$$

4. Evaluate the following:

(a) $\sum_{k=1}^4 k$

$$\sum_{k=1}^4 k = 1 + 2 + 3 + 4 = 10.$$

(b) $\sum_{k=0}^5 k^2$

$$\sum_{k=0}^5 k^2 = 0^2 + 1^2 + 2^2 + 3^2 + 4^2 + 5^2 = 0 + 1 + 4 + 9 + 16 + 25 = 55.$$

(c) $\sum_{k=1}^{10} 3$

$$\sum_{k=1}^{10} 3 = \underbrace{3 + 3 + \dots + 3}_{10} = 10 \cdot 3 = 30.$$

(d) $\sum_{k=1}^n k$

$$\begin{aligned} \sum_{k=1}^n k &= \underbrace{1 + 2 + \dots + n}_n \\ &= \underbrace{(n+1) + (n-1+2) + \dots + n/2}_{n/2} \\ &= \frac{n(n+1)}{2}. \end{aligned}$$

(e) $\sum_{k=1}^n n$

$$\sum_{k=1}^n n = \underbrace{n + n + \dots + n}_n = n^2.$$

(f) $\sum_{k=1}^n 1$

$$\sum_{k=1}^n 1 = \underbrace{1 + 1 + \dots + 1}_n = n.$$

5. The matrix C^T whose rows are the same as the respective columns of matrix C is called the transpose of C . For example,

$$C = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, C^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}.$$

(a) Let the elements of C be c_{ij} and the elements of C^T be c'_{ij} . Write a formula for C^T in terms of these elements. That is, $c'_{ij} = ?$

We simply have $c'_{ij} = c_{ji}$; the indices are swapped.

(b) Write $\begin{bmatrix} 2 & 1 & 5 \\ 4 & -2 & 0 \end{bmatrix}^T$.

We flip it over the main diagonal; since the matrix is not square, we get a matrix with different dimensions!

$$\begin{bmatrix} 2 & 1 & 5 \\ 4 & -2 & 0 \end{bmatrix}^T = \begin{bmatrix} 2 & 4 \\ 1 & -2 \\ 5 & 0 \end{bmatrix}$$

6. Fill in the blanks: Multiplying an $m \times n$ matrix by a(n) $\underline{\quad} \times k$ matrix gives a(n) $\underline{\quad} \times \underline{\quad}$ matrix.

Multiplying an $m \times n$ matrix by a(n) $\underline{n} \times k$ matrix gives a(n) $\underline{m} \times \underline{n}$ matrix.

7. Dogs can eat cats, rats, or mice; cats can eat rats or mice; rats can eat mice.

(a) Make a matrix E showing what can eat what.

Let dogs be D , cats be C , rats be R , and mice be M . Then the matrix is straightforward. Note that we put the prey on the left and predator on top because then following the matrices is going through each step in the food chain.

$$T = \begin{array}{c} \text{predator} \\ D \quad C \quad R \quad M \\ \text{prey} \begin{bmatrix} D \\ C \\ R \\ M \end{bmatrix} \end{array} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{bmatrix}.$$

Note that the diagonal is all 0s because no animal eats their own species.

(b) Draw a directed graph.

The graph is shown in Figure 1.




Figure 1: The directed graph of E .

(c) Calculate and interpret E^2 , E^3 , E^4 .

We have

$$\begin{aligned}
 E^2 &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{bmatrix} \\
 &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \end{bmatrix}.
 \end{aligned}$$

This means that there are two ways for a mouse's nutrients to find its way to a dog in two steps (namely, through a rat and through a cat). Also, there is only one way for a rat to get to a dog in two steps, and only one way for a mouse to get to a cat in two steps.

We have

$$\begin{aligned}
 E^3 = EE^2 &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \end{bmatrix} \\
 &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}.
 \end{aligned}$$

This means a mouse can get to a dog in three steps in only one way.

We have

$$\begin{aligned}
 E^4 = EE^3 &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \\
 &= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.
 \end{aligned}$$

A matrix of all zeroes! That means there are no paths which take four steps, which makes sense.

8.

Note that as given in the problem, we have

$$S = \begin{bmatrix} 5 & 1 & 0 & \frac{4}{4} & 1 & 1 & 0 & 0 \\ 5 & 1 & 1 & \frac{1}{4} & \frac{3}{4} & 0 & \frac{1}{3} & 2 \end{bmatrix};$$

$$C = [5 \quad 20 \quad 10 \quad 0 \quad 1 \quad 2 \quad 5 \quad 12].$$

(a) Unfortunately, if you try to multiply S and C as given, it won't work. Why not?

The dimensions aren't right! S is a 2×8 matrix, while C is a 1×8 matrix.

(b) What do you need to do to C so they can be multiplied? Explain the dimensions of each matrix.

We need to transpose C , since C^T is a 8×1 matrix. This lets it be multiplied by S .

(c) Once you've fixed matrix C , do the multiplication. What are the dimensions of your answer?

We do the multiplication:

$$SC^T = \begin{bmatrix} 5 & 1 & 0 & \frac{4}{4} & 1 & 1 & 0 & 0 \\ 5 & 1 & 1 & \frac{1}{4} & \frac{3}{4} & 0 & \frac{1}{3} & 2 \end{bmatrix} \begin{bmatrix} 5 \\ 20 \\ 10 \\ 0 \\ 1 \\ 2 \\ 5 \\ 12 \end{bmatrix}$$

$$= \begin{bmatrix} 48 \\ \frac{977}{12} \end{bmatrix}.$$

We end up with a 2×1 matrix. Indeed, $M_{2 \times 8} M_{8 \times 1} = M_{2 \times 1}$; in some sense the inner dimensions VANISH, ANNIHILATE, whatever you like, to leave the outer dimensions behind.

9. Matrix multiplication is not necessarily commutative, even when the dimensions of the matrices suggest it might be. How do we know? Be specific.

One way to know is to just multiply two random matrices together (chosen for computational convenience) and check for commutativity:

$$C = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}; \quad D = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

We compute CD and DC :

$$CD = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \cdot DC = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \implies CD \neq DC.$$

Oof! Matrix multiplication is not commutative.

10. Matrix multiplication is associative, though. Prove that $(PX)Y = P(XY)$ for

$$P = \begin{bmatrix} m & n \\ p & q \end{bmatrix}, \quad X = \begin{bmatrix} x & y \\ z & w \end{bmatrix}, \quad Y = \begin{bmatrix} r & s \\ t & u \end{bmatrix}.$$

With great dread, we compute $(PX)Y$ and $P(XY)$.

$$\begin{aligned}
 (PX)Y &= \left(\begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} x & y \\ z & w \end{bmatrix} \right) \begin{bmatrix} r & s \\ t & u \end{bmatrix} \\
 &= \begin{bmatrix} mx + nz & my + nw \\ px + qz & py + qw \end{bmatrix} \begin{bmatrix} r & s \\ t & u \end{bmatrix} \\
 &= \begin{bmatrix} r(mx + nz) + t(my + nw) & s(mx + nz) + u(my + nw) \\ r(px + qz) + t(py + qw) & s(px + qz) + u(py + qw) \end{bmatrix} \\
 &= \begin{bmatrix} mrx + mty + nrz + ntz & msx + muy + nsz + nuw \\ prx + pty + qrz + qtw & psx + puy + qsz + quw \end{bmatrix}. \\
 P(XY) &= \begin{bmatrix} m & n \\ p & q \end{bmatrix} \left(\begin{bmatrix} x & y \\ z & w \end{bmatrix} \begin{bmatrix} r & s \\ t & u \end{bmatrix} \right) \\
 &= \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} xr + yt & xs + yu \\ zr + wt & zs + wu \end{bmatrix} \\
 &= \begin{bmatrix} m(xr + yt) + n(zr + wt) & m(xs + yu) + n(zs + wu) \\ p(xr + yt) + q(zr + wt) & p(xs + yu) + q(zs + wu) \end{bmatrix} \\
 &= \begin{bmatrix} mrx + mty + nrz + ntz & msx + muy + nsz + nuw \\ prx + pty + qrz + qtw & psx + puy + qsz + quw \end{bmatrix}.
 \end{aligned}$$

They are equal!

Note that this doesn't prove it's associative for all qualifying¹⁸ matrices because we've only shown it for 2×2 matrices.

11. Prove that matrix multiplication is distributive: $P(X + Y) = PX + PY$.

This isn't as bad.

$$\begin{aligned}
 P(X + Y) &= \begin{bmatrix} m & n \\ p & q \end{bmatrix} \left(\begin{bmatrix} x & y \\ z & w \end{bmatrix} + \begin{bmatrix} r & s \\ t & u \end{bmatrix} \right) \\
 &= \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} x + r & y + s \\ z + t & w + u \end{bmatrix} \\
 &= \begin{bmatrix} m(x + r) + n(z + t) & m(y + s) + n(w + u) \\ p(x + r) + q(z + t) & p(y + s) + q(w + u) \end{bmatrix} \\
 &= \begin{bmatrix} (mx + nz) + (mr + nt) & (my + nw) + (ms + nu) \\ (px + qz) + (pr + qt) & (py + qw) + (ps + qu) \end{bmatrix} \\
 &= \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} x & y \\ z & w \end{bmatrix} + \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} r & s \\ t & u \end{bmatrix} \\
 &= PX + PY.
 \end{aligned}$$

Indeed, they are equal!

12. When does $PX = XP$? Don't worry if you get some messy equations in your answer.

Let's try it.

$$\begin{aligned}
 PX &= \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} x & y \\ z & w \end{bmatrix} \\
 &= \begin{bmatrix} mx + nz & my + nw \\ px + qz & py + qw \end{bmatrix} \\
 XP &= \begin{bmatrix} x & y \\ z & w \end{bmatrix} \begin{bmatrix} m & n \\ p & q \end{bmatrix} \\
 &= \begin{bmatrix} xm + yp & xn + yq \\ zm + wp & zn + wq \end{bmatrix}
 \end{aligned}$$

¹⁸In terms of dimension.

Equating terms, we get

$$\begin{cases} mx + nz = xm + yp \\ my + nw = xn + yq \\ px + qz = zm + wp \\ py + qw = zn + wq \end{cases}$$

$$\Rightarrow \begin{cases} nz = yp \\ my + nw = xn + yq \\ px + qz = zm + wp \\ py = zn \end{cases}$$

13. Cook's Seafood Restaurant in Menlo Park sells fish and chips. The Captain's order is two pieces of fish and one order of chips, while the Regular order is one piece of fish and one order of chips.

(a) Write a matrix representing these facts, with clear labels on your rows and columns.

Let Captain = C and Regular = R . Then the matrix is:

$$M = \begin{array}{c} \text{fish} \quad \text{chips} \\ C \quad \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \\ R \end{array}$$

I'm writing it this way rather than the transpose so that the order of the matrices in the next problem is the same as the problems appear. Otherwise, you'd have to flip the order of multiplication (remember, it's not commutative!).

(b) The restaurant management estimates their cost at 0.75 for each piece of fish and 0.50 for each order of chips. Represent this as a matrix, then use matrix multiplication to calculate the cost of the two possible orders.

$$N = \begin{array}{c} \text{cost (\$)} \\ \text{fish} \quad \begin{bmatrix} 0.75 \\ 0.50 \end{bmatrix} \\ \text{chips} \end{array}$$

Now we just multiply the matrices:

$$\begin{aligned} MN &= \begin{array}{c} \text{fish} \quad \text{chips} \\ C \quad \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix} \\ R \end{array} \begin{array}{c} \text{cost (\$)} \\ \text{fish} \quad \begin{bmatrix} 0.75 \\ 0.50 \end{bmatrix} \\ \text{chips} \end{array} \\ &= \begin{array}{c} \text{cost (\$)} \\ C \quad \begin{bmatrix} 2 \\ 1.25 \end{bmatrix} \\ R \end{array} \end{aligned}$$

Thus, the cost of a Captain's order is \$2 and the cost of a Regular order is \$1.25 (for the restaurant).

(c) For a party, Cook's provides 10 Captain's orders and 5 Regular orders. Write this as a matrix and use matrix multiplication to find how many pieces of fish and orders of chips are provided.

We want to multiply this matrix by M and get a 2×1 or 1×2 matrix of fish and chips. Thus, we choose the 1×2 matrix

$$P = \begin{matrix} C & R \\ 10 & 5 \end{matrix},$$

so that the product is simply

$$PM = \begin{matrix} C & R \\ 10 & 5 \end{matrix} \begin{matrix} C & R \\ \text{fish} & \text{chips} \\ \left[\begin{matrix} 2 & 1 \\ 1 & 1 \end{matrix} \right] \end{matrix} = \begin{matrix} \text{fish} & \text{chips} \\ 25 & 15 \end{matrix}.$$

Thus, 25 pieces of fish and 15 orders of chips are provided.

(d) Now use matrix multiplication to find out the cost of the party.

We need to multiply PM by N to get a 1×1 matrix:

$$PMN = \begin{matrix} \text{fish} & \text{chips} \\ 25 & 15 \end{matrix} \begin{matrix} \text{fish} & \text{chips} \\ \left[\begin{matrix} 0.75 \\ 0.50 \end{matrix} \right] \end{matrix} = [26.25].$$

Thus, the party costs \$26.25 for the restaurant.

14. We will find coefficient matrices to be particularly useful for solving systems of linear equations.

For instance,

$$\begin{cases} 3x + 4y = 5 \\ 6x + 4y = 8 \end{cases} \leftrightarrow \begin{bmatrix} 3 & 4 \\ 6 & 7 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 8 \end{bmatrix}.$$

Rewrite

$$\begin{cases} 2x + 3y + 4z = 5 \\ 5x - 4y + 2z = 2 \\ x + 2y = 7 \end{cases}$$

as a matrix equation in this way.

We rewrite the last equation as $x + 2y + 0z = 7$ and proceed to tabulate the coefficients:

$$\underbrace{\begin{bmatrix} 2 & 3 & 4 \\ 5 & -4 & 2 \\ 1 & 2 & 0 \end{bmatrix}}_M \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 2 \\ 7 \end{bmatrix}.$$

15.

(a) What is the transpose of the 3×3 matrix M from the previous problem?

M is shown in the previous problem above. We flip it, obtaining

$$M^T = \begin{bmatrix} 2 & 5 & 1 \\ 3 & -4 & 2 \\ 4 & 2 & 0 \end{bmatrix}.$$

(b) Use M^T to rewrite the system in the previous problem.

At first, one might try flipping the order of the column matrix to say z, y, x (from top to bottom). But this doesn't work.

Thinking in terms of what columns and rows mean, we can label them in the original equation from the previous problem:

$$\begin{array}{l} \text{Eq. 1} \\ \text{Eq. 2} \\ \text{Eq. 3} \end{array} \begin{array}{l} x \\ y \\ z \end{array} \begin{array}{l} 2 \\ 5 \\ 1 \end{array} \begin{array}{l} 3 \\ -4 \\ 2 \end{array} \begin{array}{l} 4 \\ 2 \\ 0 \end{array} \begin{array}{l} x \\ y \\ z \end{array} \begin{array}{l} [x] \\ [y] \\ [z] \end{array} = \begin{array}{l} \text{Eq. 1} \\ \text{Eq. 2} \\ \text{Eq. 3} \end{array} \begin{array}{l} [5] \\ [2] \\ [7] \end{array}.$$

Doing the same for M^T , we get

$$M^T = \begin{array}{l} x \\ y \\ z \end{array} \begin{array}{l} \text{Eq. 1} \\ \text{Eq. 2} \\ \text{Eq. 3} \end{array} \begin{array}{l} 2 \\ 3 \\ 4 \end{array} \begin{array}{l} 5 \\ -4 \\ 2 \end{array} \begin{array}{l} 1 \\ 2 \\ 0 \end{array} \end{array}.$$

We realize that to get a matrix containing the values of Equations 1-3, we need to left-multiply M^T by a 1×3 matrix $[x \ y \ z]$:

$$[x \ y \ z] \begin{array}{l} 2 \\ 3 \\ 4 \end{array} \begin{array}{l} 5 \\ -4 \\ 2 \end{array} \begin{array}{l} 1 \\ 2 \\ 0 \end{array} = [5 \ 2 \ 7].$$

This looks quite similar to the previous problem! In symbols, if $A = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ and $B = \begin{bmatrix} 5 \\ 2 \\ 7 \end{bmatrix}$, we have

$$\underbrace{MA = B}_{\text{prev. prob.}} \leftrightarrow A^T M^T = B^T.$$

(c) What is the transpose of the transpose matrix, $(M^T)^T$?

Since we're just reflecting over the diagonal twice, we have $(M^T)^T = M$.

This brings up an interesting fact, thinking back to the last subproblem. In general for any matrices P, Q, R , we have

$$PQ = R \leftrightarrow Q^T P^T = R^T.$$

Taking the transpose of both sides of the right equation, we get

$$(Q^T P^T)^T = (R^T)^T = R.$$

Equating this with the left equation, we get

$$PQ = (Q^T P^T)^T.$$

Succulent!